



INFLUENCE OF APPLIED NITROGEN FERTILIZER ON THE BIOACCUMULATION OF MICRONUTRIENTS, ANTI-NUTRIENTS AND TOXIC SUBSTANCES IN *TELFAIRIA OCCIDENTALIS* (FLUTED PUMPKIN).

Amanabo Musa<sup>\*1,2</sup> and Emmanuel O. Ogbadoyi<sup>2</sup>

<sup>1</sup> Present address: Department of Biochemistry, Ibrahim Badamasi Babangida University, Lapai, Niger State, Nigeria

<sup>2</sup> Department of Biochemistry, Federal University of Technology, Minna, Nigeria

\*Corresponding author: Dr. Amanabo Musa is a Senior Lecturer and Head of Department of Biochemistry at Ibrahim Badamasi Babangida University, Lapai, Niger State, Nigeria E mail address:

[musaamanabo@gmail.com](mailto:musaamanabo@gmail.com) Tel: + 234 (0)803 488 4031

**ABSTRACT:** The chemical contents and compositions of plants are influenced by the physical and chemical properties of the soils on which they were grown. It is on this grounds that pot experiment was carried out to investigate the effect of soil nitrogen levels on antinutrients (soluble and total oxalates), toxic substances (cyanide and nitrate) and some micronutrients namely, vitamin C,  $\beta$ -carotene (precursor of vitamin A) and mineral elements (Fe, Mg, Zn, Cu, Ca, Na and K) in *Telfairia occidentalis*. The leaves of *Telfairia occidentalis* were harvested at market maturity (vegetative phase) and fruiting (reproductive phase) and they were subjected to chemical analysis. Results obtained showed that the applied nitrogen fertilizer significantly ( $p < 0.05$ ) increased the cyanide and nitrate concentrations in *Telfairia occidentalis* in both phases of plant development. The soluble and total oxalate concentrations in the vegetable were not significantly affected by the applied nitrogen in both stages of its development. Similarly, the increasing effect of nitrogen fertilizer on  $\beta$ -carotene content in *Telfairia occidentalis* was not significant at vegetative phase, but its concentration was increased significantly ( $p < 0.05$ ) at reproductive phase. Vitamin C concentration in *Telfairia occidentalis* was reduced significantly ( $p < 0.05$ ) with the applied nitrogen fertilizer irrespective of the phase of its development. The concentrations of Mg, Ca, Na and K were not significantly affected by the applied nitrogen fertilizer, however, the levels of Zn and Cu in *Telfairia occidentalis* were increased significantly at market maturity and fruiting, respectively. While the concentration of Fe in the vegetable increased significantly at both stages of plant development. The result concludes that the application of nitrogen fertilizer for the cultivation of *Telfairia occidentalis* increases the concentration of plant toxins and reduces the nutritional quality of the vegetable.

**Keywords:** *Telfairia occidentalis*, soil nitrogen levels, oxalates, nitrate, cyanide, micronutrients.

## INTRODUCTION

Fluted pumpkin (*Telfairia occidentalis*) popularly called “ugwu” by Igbos, is a creeping vegetative shrub that spread low across the ground with large lobed leaf, and long twisting tendrils (Sydenham, 1985; Horsfall and Spiff, 2005; Christian, 2006; Ojiako and Igwe, 2008). Nkang *et al.* (2003) reported that this vegetable is of commercial importance grown across the low land humid tropic in West Africa (Nigeria, Ghana and Sierra Leone being the major producers). Fluted pumpkin prefers a loose, friable ample humus and shaded position. Nitrogen is essential for adequate vegetation and should ideally be given in the form of manure (Schippers, 2000). The leaves and seeds are widely eaten as they are good sources of minerals (potassium, magnesium, sodium, phosphorus and iron),  $\beta$ -carotene, vitamins, fibres, fats (Schipper, 2000; Nkang *et al.*, 2003; Christian, 2006; Ogbadoyi *et al.*, 2011).

Harvesting of fluted pumpkin takes place 120 - 150 days after planting. The seed contains 13% of oil (Okoli and Nyanayo, 1988) and is used for cooking (Horsefall and Spiff, 2005).

*Telfairia occidentalis* like other leafy vegetables also contains appreciable levels of some antinutrients and toxic substance which has been shown to have negative effect on animal and human health at high concentrations (Morton, 1987; Ojokoh *et al.*, 2002; Ogbadoyi *et al.*, 2011; Musa *et al.*, 2011). Ekpedema *et al.* (2000) reported the presence of antinutritional factors (oxalate and phytate) and other toxic substances in the leaves and seeds of *Telfairia occidentalis*.

Various cultural practices like optimum N, P and K fertilizer levels required for adequate vegetative growth and yield in *Telfairia occidentalis* has been established. However, the effect of these optimal fertilizer levels on the nutrients, antinutrients and toxic substances has not been determined. The present study is therefore, designed to investigate the influence of soil fertility in terms of N levels on the accumulation of some micronutrients and phytotoxins in the leaves of *Telfairia occidentalis*.

## **MATERIALS AND METHODS**

### **The study area**

The pot experiment was conducted in the nursery of the School of Agriculture and Agricultural Technology, Federal University of Technology, Minna, Niger State of Nigeria.

Niger State has a Savannah climate characterized by maritime. The geographical location of Minna is at longitude 9° 40'N and latitude 6° 30'E. Minna lies in the Southern Guinea Savannah zone of Nigeria and has a sub - humid semi arid tropical climate. The raining season is between April and October. About 90% of the total rainfall occurs between the month of June and September. The mean annual rainfall is in the range of 1200 – 1300 mm. The temperature of this zone rarely falls below 22°C with peaks temperature of 40°C in February /March and 30°C in November /December. Wet season average temperature is about 29°C. The Dry season occurs between November and March while harmattan which is characterised by dry air is between November and February (Osunde and Alkassoun, 1998).

### **Soil sampling and analysis**

The soil used in this study was collected from Minna. The soil has been classified as Inceptisol (FDALR, 1985). The bulked sample was collected during the dry season from the field which has been under fallows for about four years. The bulked soil sample was passed through 2 mm sieve. Sub-sample of the soil was subjected to routine soil analysis using procedure described by Juo (1979). The soil particle sizes were analyzed using hydrometer method; pH was determined potentiometrically in the water and 0.01M CaCl<sub>2</sub> solution in a 1: 2 soil/ liquid using a glass electrode pH meter and organic carbon by Walkey-Black method (Juo, 1979). Exchange acidity (E.A H<sup>+</sup> and Al<sup>3+</sup>) was determined by titration method (Juo, 1979). Exchangeable Ca, Mg, K and Na were leached from the soil sample with neutral 1N NH<sub>4</sub>OA solution. Sodium and potassium were determined by flame emission spectrophotometry while Mg and Ca were determined by E.D.T.A versenate titration method (Juo, 1979). Total nitrogen was estimated by Macrokjedal procedure and available phosphorus by Bray No 1 method (Juo, 1979).

### **Sources of Seeds**

The seeds of fluted pumpkin (*Telfairia occidentalis*) were obtained from Schools of Agriculture and Agricultural Technology's Nursery of Federal University of Technology, Minna.

### **Planting, experimental design and nursery management**

Exactly two seeds of *Telfairia occidentalis* were planted in a polythene bag filled with 20.00 kg of top soil and after germination the seedlings were thinned to one plant per pot. The factorial design was adopted to determine the effect nitrogen fertilizer at vegetative and reproductive phases in *Telfairia occidentalis*.

Each treatment had 10 pots replicated three times. This gave a total of 60 pots for *Telfairia occidentalis*. The seedlings were watered twice daily (morning and evening) using watering can and weeded regularly. The experimental area and the surroundings were kept clean to prevent harbouring of pest. The pots were lifted from time to time to prevent the roots of the plants from growing out of the container. Insects were controlled using Sherpa plus (Saro Agro Sciences) four weeks after planting at the rate of 5 ml per 5 litres of water.

### **Fertilizer treatment**

Fertilizer treatment for the vegetable was N fertilization at two levels. The first level was control (no N fertilization) and the second level was N fertilization. Basal application of P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O in the form of single super phosphate and murate of potash was done. The details of fertilizer treatments are as follow:

F<sub>1</sub> (control): 0N, 30mg P<sub>2</sub>O<sub>5</sub>/kg soil and 22mg K<sub>2</sub>O/kg soil

F<sub>2</sub>: 30mgN/kg soil, 30mg P<sub>2</sub>O<sub>5</sub>/kg soil and 22mg K<sub>2</sub>O/kg soil

### Harvesting

The leaves of *Telfairia occidentalis* grown in pot experiment in control and nitrogen treated soil were harvested at market maturity and fruiting.

### Sample analysis

The soluble and total oxalates concentration in the leaves of *Telfairia occidentalis* were determined by titrimetric method of Oke (1966). Nitrate concentration in the samples was determined by the colourimetric method (Sjoberg and Alanka, 1994). Alkaline picrate method was used to analyse the cyanide concentration (Ikediobi *et al.*, 1980). The mineral elements (Fe, Mg, Zn, Cu, Ca, Na and K) in samples were determined according to the method of Ezeonu *et al.* (2002). The ascorbic acid concentration was determined by 2, 6-dichlorophenol indophenols method (Eleri and Hughes, 1983) while the estimation of  $\beta$ -carotene was done by ethanol and petroleum ether extraction method (Musa *et al.*, 2010).

### Statistical analysis

T-test was used to determine the effect of soil fertility using two levels of nitrogen fertilizer on the level of the parameters under investigation.

## RESULTS

### Physical and chemical properties of soil

Result of analysis of the soil used for pot experiment is presented in Table 1. The texture class of the soil is sandy loam indicating that the water holding capacity is moderate. The organic matter content, total nitrogen and available phosphorus are low. Sodium and calcium contents are moderate while magnesium and potassium contents are high. The CEC (cation exchange capacity) is moderate while base saturation percentage is high. Soil pH indicates that the soil is slightly acidic (FAO, 1984; Black, 1985; FDALR, 1985).

### Effect of nitrogen fertilizer on antinutrients and vitamins content

The results obtained from the determination of cyanide concentration in *Telfairia occidentalis* indicated that the applied nitrogen fertilizer significantly ( $p < 0.05$ ) increased the cyanide concentration in the vegetable irrespective of the stage of plant development. The mean values of cyanide in control and nitrogen fertilized *Telfairia occidentalis* at vegetative phase were  $438.00 \pm 37.00$  mg/kg and  $699.00 \pm 48.00$  mg/kg while the values obtained at reproductive phase were  $771.00 \pm 21.00$  mg/kg and  $885.00 \pm 37.00$  mg/kg (Table 2).

The applied nitrogen fertilizer also increased nitrate concentration significantly ( $p < 0.05$ ) in *Telfairia occidentalis* at vegetative and reproductive phases. The mean values of nitrate in the control and nitrogen applied vegetable at market maturity were  $550.00 \pm 95.00$  mg/kg and  $696.00 \pm 117.00$  mg/kg while the corresponding values obtained at fruiting were  $34.90.00 \pm 5.1.00$  mg/kg and  $45.10 \pm 15.00$  mg/kg (Table 2).

Application of nitrogen fertilizer had no significant effect ( $p > 0.05$ ) on soluble and total oxalate concentrations at both stages of the plant development. (Table 2).

The applied nitrogen fertilizer significantly ( $p < 0.05$ ) elevated the  $\beta$ -carotene concentration in *Telfairia occidentalis* at market maturity and had no significant ( $p > 0.05$ ) effect on its concentration at fruiting. The amount of  $\beta$ -carotene concentration in control and nitrogen treated *Telfairia occidentalis* at market maturity were  $15501.00 \pm 591.00$   $\mu$ g/100 g and  $17600 \pm 1009$   $\mu$ g/100 g while the corresponding values at fruiting were  $10381.00 \pm 395.79$  g/100g and  $12150.00 \pm 276.00$   $\mu$ g/100 g (Table 2).

The vitamin C concentration in control and nitrogen applied *Telfairia occidentalis* at market maturity were  $208.40 \pm 7.50$  mg/100 g and  $191.60 \pm 13.00$  mg/100g while the values obtained at fruiting were  $224.60 \pm 11.00$  mg/100 g and  $186.30 \pm 11.00$  mg/100 g. The results showed that the application of nitrogen fertilizer significantly decreased ( $p < 0.05$ ) the vitamin C concentration in *Telfairia occidentalis* at both stages of plant development (Table 2).

### Effect of nitrogen fertilizer on mineral elements content

The application of nitrogen fertilizer significantly ( $p < 0.05$ ) increased Fe concentration in *Telfairia occidentalis* irrespective of the phase of its development. The mean value of Fe concentrations in nitrogen applied vegetable at market maturity ( $13.63 \pm 1.30$  mg/kg) and fruiting ( $28.19 \pm 2.40$  mg/kg) were significantly higher when compared with the values of controls ( $10.76 \pm 0.92$  mg/kg and  $17.84 \pm 2.40$  mg/kg respectively), (Table 3).

**Table 1: Some Physical and Chemical Properties of the Soil (0 – 20cm) Used for Pot Experiment**

Parameters	Values
Sand (%)	74.40
Silt (%)	18.00
Clay (%)	7.60
pH (in H <sub>2</sub> O)	6.51
pH (in 0.1M C <sub>a</sub> Cl <sub>2</sub> )	5.25
Organic Carbon (%)	0.83
Organic Matter (%)	1.43
Total nitrogen (%)	0.05
Available phosphorus (mg/kg)	6.69
K (cmol/kg)	0.92
Na (cmol/kg)	0.68
Mg (cmol/kg)	4.80
Ca (cmol/kg)	8.00
E. A (H <sup>+</sup> +AL <sup>3+</sup> )(cmol/kg)	1.50
CEC (cmol/kg)	15.90
Base saturation (%)	90.57
Texture class	sandy loam

\*Values represent means of triplicate determinations.

The applied nitrogen fertilizer had no significant effect on Mg, Ca, Na and K concentrations of the vegetable at both stages of plant development (Table 3).

There was a significant ( $p < 0.05$ ) increase in Zn concentration at market maturity with nitrogen fertilization while no significant variation was observed at fruiting (Table 3).

**Table 2: Effect nitrogen fertilizer on antinutrients and vitamins content in *Telfairia occidentalis***

Antinutrients and vitamins analysed at market maturity and fruiting stage	Nitrogen levels	
	Control (No. nitrogen applied)	Nitrogen applied
Cyanide at market maturity (mg/kg DW)	438.00 ± 37.00 <sup>a</sup>	699.00 ± 48.00 <sup>b</sup>
Cyanide at fruiting (mg/kg DW)	771.00 ± 21.00 <sup>a</sup>	885.00 ± 37.00 <sup>b</sup>
Nitrate at market maturity (mg/kg DW)	550.00 ± 95.00 <sup>a</sup>	696.00 ± 117.00 <sup>b</sup>
Nitrate at fruiting (mg/kg DW)	34.90 ± 5.10 <sup>a</sup>	45.90 ± 15.00 <sup>b</sup>
Soluble at market maturity (g/100 g DW)	1.64 ± 0.06 <sup>a</sup>	1.82 ± 0.14 <sup>a</sup>
Soluble at fruiting (g/100 g DW)	1.65 ± 0.09 <sup>a</sup>	2.03 ± 0.06 <sup>a</sup>
Total oxalate at market maturity (g/100 g DW)	2.18 ± 0.07 <sup>a</sup>	2.01 ± 0.16 <sup>a</sup>
Total oxalate at fruiting (g/100 g DW)	3.20 ± 0.09 <sup>a</sup>	2.82 ± 0.04 <sup>a</sup>
β-carotene at market maturity (μg/100 g FW)	15501.00 ± 591.00 <sup>a</sup>	17600.30 ± 100.00 <sup>b</sup>
β-carotene at fruiting (μg/100 g FW)	10381.00 ± 395.79 <sup>a</sup>	12150.00 ± 276.00 <sup>a</sup>
Vitamin C at market maturity (mg/100 g FW)	208.40 ± 7.50 <sup>b</sup>	191.60 ± 13.00 <sup>a</sup>
Vitamin C at fruiting (mg/100 g FW)	224.60 ± 11.00 <sup>b</sup>	186.30 ± 11.00 <sup>a</sup>

DW = Dry weight, FW = Fresh weight, values represent means of nine determinations. Row mean values carrying the same superscripts do not differ significantly from each other ( $P > 0.05$ ).

Application of nitrogen fertilizer had no significant effect on the Cu concentration in *Telfairia occidentalis* at market maturity, however, at fruiting, the concentration of the mineral element increased significantly ( $p < 0.05$ ). The mean values in the control and nitrogen treated vegetable at market maturity were  $3.67 \pm 0.12$  mg/kg and  $2.66 \pm 0.53$  mg/kg while at fruiting the values obtained were  $1.30 \pm 0.51$  mg/kg and  $8.78 \pm 0.72$  mg/kg (Table 3)

**Table 3: Effect of nitrogen fertilizer on mineral contents in *Telfairia occidentalis***

Mineral analysed at market maturity and fruiting stages.	Nitrogen levels	
	Control (No nitrogen applied)	Nitrogen applied
Fe at market maturity (mg/kg)	10.76 ± 0.92 <sup>a</sup>	13.63 ± 1.30 <sup>b</sup>
Fe at fruiting (mg/kg)	17.84 ± 2.40 <sup>a</sup>	28.19 ± 2.40 <sup>b</sup>
Mg at market maturity (mg/kg)	19.90 ± 1.00 <sup>a</sup>	20.34 ± 0.80 <sup>a</sup>
Mg at fruiting (mg/kg)	23.13 ± 2.24 <sup>a</sup>	23.59 ± 3.06 <sup>a</sup>
Zn at market maturity (mg/kg)	0.03 ± 0.01 <sup>a</sup>	0.05 ± 0.01 <sup>b</sup>
Zn at fruiting (mg/kg)	0.01 ± 0.01 <sup>a</sup>	0.02 ± 0.01 <sup>a</sup>
Cu at market maturity (mg/kg)	3.67 ± 0.12 <sup>a</sup>	2.66 ± 0.53 <sup>a</sup>
Cu at fruiting (mg/kg)	1.30 ± 0.51 <sup>a</sup>	8.78 ± 0.72 <sup>b</sup>
Ca at market maturity (mg/kg)	17.76 ± 3.20 <sup>a</sup>	16.27 ± 2.40 <sup>a</sup>
Ca at fruiting (mg/kg)	16.27 ± 2.00 <sup>a</sup>	14.95 ± 1.40 <sup>a</sup>
Na at market maturity (mg/kg)	5.40 ± 0.54 <sup>a</sup>	5.21 ± 0.16 <sup>a</sup>
Na at fruiting (mg/kg)	4.59 ± 0.38 <sup>a</sup>	3.71 ± 0.29 <sup>a</sup>
K at market maturity (mg/kg)	117.40 ± 3.40 <sup>a</sup>	112.90 ± 7.40 <sup>a</sup>
K at fruiting (mg/kg)	62.47 ± 3.00 <sup>a</sup>	67.40 ± 3.80 <sup>a</sup>

Values represent means of nine determinations. Row mean values carrying the same superscripts do not differ significantly from each other ( $P > 0.05$ ).

## DISCUSSION

The significantly higher cyanide, nitrate and  $\beta$ -carotene concentrations in the *Telfairia occidentalis* grown on soil treated with nitrogen compared with the control is in accordance with the report of the following authors (Kriedenman, 1964; Jones and Ford, 1972; Richard, 1991; Yang, 1992; Chweya, 1993; Oladele *et al.*, 1997; Muramoto, 1999; Peter and Birger 2002; Waclaw and Stefan, 2004; Kansal *et al.* (2005); Carmen *et al.*, 2007; Anjana and Muhammad, 2006; Anjana *et al.*, 2007; Rolinda and Ma, 2008; Musa *et al.*, 2010; Musa *et al.*, 2011). Virginia (2001) stressed that plants require nitrogen for normal growth and protein synthesis however, if nitrogen is applied in excess of what the plant requires for protein formation, the excess is accumulated as nitrate and stored predominantly in the green leafy part of the plant. Peter and Birger (2002) stated that the increased in cyanide concentration following the application of nitrogen fertilizer is that the applied nitrogen stimulates the enzymatic conversion of tyrosine to p - hydroxymandelonitrile which ultimately lead to increase in the biosynthesis of cyanogenic glycoside. The higher concentration of these phytotoxins (cyanide and nitrate) is enough warning signal why we must take adequate precautions when growing our vegetables with synthetic fertilizers because of the negative effect of these plant toxins in our health. For instance, cyanide is a respiratory poison while nitrate is responsible for cancer and methaemoglobinemia in human and other animals at higher concentrations. This study thus suggest that the attendant health problems associated with high levels of cyanide and nitrate can be reduced or minimised by careful application of nitrogen fertilizer at levels much lower the current dose or possible its use is completely avoided (Musa *et al.*, 2010; Musa *et al.* 2011).

Similarly, the increase in  $\beta$ -carotene may be probably due to elevation in the content and activity of chlorophyll and associated light absorbing pigments (including carotenoids) following the application of nitrogen fertilizer (Taiz and Zeiger, 2002; Havling *et al.*, 2006). Though the concentration of the provitamin A that helps to maintain good sight and act as a powerful antioxidant is increased in *Telfairia occidentalis* with application of synthetic nitrogen fertilizer, its concentration in controls can provide enough of vitamin A to meet the adult recommended daily allowance of 900  $\mu\text{g}$ .

Thus this increase in  $\beta$ -carotene alone is not a good reason to encourage the application of nitrogen fertilizer for the cultivation of this vegetable, bearing in mind the attendant health problems associated with high concentrations of cyanide and nitrate.

The insignificant effect of nitrogen fertilizer on soluble and total oxalates concentration in *Telfairia occidentalis* observed in this study is invariance with the findings of Chweya (1993) and Singh (2005). These authors reported that nitrogen fertilizers decreased the oxalate content in *Cleome gynandra* and *Chemopodium amaranticolor*, respectively. Singh (2005) attributed the decrease due to the fact that nitrogen and anions generally reduced the levels of the antinutrients since they compete with oxalate for cations and depressed the oxalate synthesis. However, the observed variation in this work from those of the previous works could suggest that the effect of nitrogen fertilizer on oxalate concentration may vary from cultivar to cultivars and other environmental factors. However, this variation is subject to further study.

The significantly lower concentration of vitamin C in *Telfairia occidentalis* following the application of inorganic nitrogen fertilizer is in harmony with the finding of the following researchers (Chweya, 1993; Virginia, 2001; Mozafar, 2005; Musa *et al.*, 2010; Musa *et al.*, 2011) to the effect that the application of nitrogen fertilizer significantly reduced the leaf vitamin C content in the vegetables. Virginia (2001) further stressed that the observed decrease in vitamin C concentration results from the increase in protein production and decrease in carbohydrate production following the application of nitrogen fertilizer. Since vitamin C is formed from carbohydrates, its synthesis also reduces. Considering the pivotal of this water soluble vitamin in human health, application nitrogen fertilizer in growing *Telfairia occidentalis* for human consumption as leafy vegetable should not be encouraged as this practice decrease the vitamin concentration significantly. Thus consumers may be denied of the full derivable nutritional benefits from the vegetable.

The significantly higher levels of Fe, Zn, and Cu in nitrogen applied *Telfairia occidentalis* agreed with the findings of other workers. Ojeniyi and Adeniyi (1999), Tarfa *et al.* (2001), Kansal *et al.* (2005), Safaa and Abd El Fattah (2007) reported that nitrogen fertilizer increased the mineral content in maize leaf, *Spinacea oleracea* and lettuce plants respectively. Safaa and Abd El Fattah (2007) attributed the increase in the mineral content (especially Fe) to increase in the levels of chlorophyll following the application of nitrogen fertilizer. The observations by these authors however, disagreed with the results of Chweya (1993) who reported that application of nitrogen fertilizer to *Gynandropsis gynandra* decreases the Fe content of the vegetable and has no significant effect on the Ca and Na content. The observed variations in the mineral content in this vegetable from the report of other authors following the application of nitrogen fertilizer could be as a result of differences in cultivars and environmental factors (Chweya and Nameus, 1997; Grazyna and Waldemar, 1999; Bolanle *et al.*, 2004; Singh, 2005; Aliyu and Morufu, 2006; Lisiewska *et al.*, 2006; Rickman *et al.*, 2007). This increased concentration of Zn and Cu in the *Telfairia occidentalis* treated with nitrogen fertilizer has also no much of nutritional advantage of the minerals than the controls. This is because concentration of these mineral elements in *Telfairia occidentalis* in both soil nitrogen levels were lower than the recommended daily allowance of 10 – 15 mg/kg and 1.3 – 3.0 mg/kg for Zn and Cu respectively, (Titz *et al.*, 1994) if 100g of the vegetable were consumed. Thus considering important roles of these mineral elements in metabolism, majorly as cofactors for many enzymatic reactions, supplementation of the elements from other sources in either case will be apparently necessary. Even with the Fe that the concentration increased significantly with applied nitrogen fertilizer, at both market maturity and fruiting, it was only at fruiting that the concentration of Fe can meet the recommended daily allowance of 18 mg/day (Titz *et al.*, 1994). However, Musa *et al.* 2011 had reported unsafe high levels of some phytotoxins in *Telfairia occidentalis* at reproductive phase and advised the consumption of the vegetable at vegetative phase rather than at reproductive phase.

## CONCLUSION

This study concludes that growing of *Telfairia occidentalis* with application of nitrogen fertilizer increase the concentrations of phytotoxins (particularly cyanide and nitrate) and reduce the nutritional quality of the vegetable. The study therefore suggest that application of nitrogen fertilizer for growing *Telfairia occidentalis* should be avoided, however, where this practice is inevitable it should be done with cautions because of the health problems associated with high ingestion of these plant toxins.

## ACKNOWLEDGEMENTS

Amanabo Musa acknowledges the contribution of his late junior brother Master Anthony Alhaji Musa for his moral and financial contribution towards the success of this research work. May his gentle soul rest in perfect peace (Amen).

## REFERENCES

- Aliyu, H.M and Morufu, A.I (2006). Proximate analysis of some leafy vegetables (Roselle, jute and bitter leaf). Int J Foods Agric. Res., 3 (1): 194 - 198.
- Anjana, S.U and Muhammad, I (2006). Nitrate accumulation in plants, factors affecting the process, and human health implications. A review. Agro. Sustainable Dev., 101: 1051 – 1060.
- Anjana, S.U., Muhammed, I and Abrol, Y.P (2007). Are nitrate concentrations in leafy vegetables within safe limits? Curr Sci., 92(3): 355 – 360.
- Black, C.A (1985). Methods of soil chemical analysis and microbiological properties. The Americal Society of Agronomy. Inc.Maidson, Wisconsin, USA. Pp 1592.
- Bolanle, A.O, Olumuyiwa, S.F., Onome, U., Bridget, O.O., A dewale, O and Steve, R.A.A (2004). The Effect of seasoning salts and local condiments on mineral availability from two Nigeria vegetables. Pak. J. Nutri., 3 (3): 146 – 153.
- Carmen, W., Angelita, D.S and Henrique, V.P (2007). Antinutrients in the cassava (*Manihot esculenta crantz*) leaf powder at three ages of the plant. Ciencia e Tecnologia de Alimentos. <http://www.Scielo.br/SciELO.php? Script = Sciarttext & pid = S 0101- 20612007000100019 & Ing = e --->
- Christian, A (2006). Studies of selected Physico chemical properties of fluted pumpkin (*Telfaria occidentalis* Hook f) Seed oil and tropical almond (*Terminalia catappia* l) Seed oil. Pak. J. Nutr., 5 (4): 306 – 307.
- Chweya, J.A (1993). Genetic enhancement of indigenous vegetables in Kenya; Field and laboratory experience report. Kengo, Nairobi. <http://www.ipgri.cgiar. Org/publications/HTMLpublications/500/ch12.htm>
- Chweya, J.A and Nameus, A.M (1997). Cats whiskers (*Cleome gynandra* l): Promoting the conservation and use of underutilized and neglected crops. II. Institute of plant genetics and crop plant research. Gatersleben / International plant Genetic Resources Institute, Rome, Italy. Pp 18 - 21.
- Ekpedema, U.A., Bassy, A.N and Ekaete, UE (2000). Minerals and antinutrients in fluted pumpkin (*Telfairia occidentalis*). Food Chem., 70 (2): 235 – 240.
- Eleri, T and Hughes, R.E (1983). Foliar ascorbic acid in some angiosperms. Phytochem. 2 (11): 2493.
- Ezeonu, F.C., Musa, A., Stanly, C.D., Oswald, C.C (2002). Iron and zinc status in soils, water and stable food cultivars in Itakpe, Kogi state of Nigeria. Environmentalist. 22; 237-240.
- FAO (1984). Fertilizers and plant nutrition. Bulletin No.9. Food and Agricultural Orgaanization. Romme.
- FDALR (1985). The reconnaissance soil survey of Niger State. Soil report. Federal Department of Agricultural Land Resources.
- Grazyna, J. and Waldemar, K (1999). Content of selected mineral compounds, nitrates III and V. and oxalates in spinach (*Spinacia oleracea* L.) and new zeal and spinach (*Tetragonia espansamurr.*) from spring and autumn growing seasons. Electronic Journal of Polish Agricultural Universities, Food Sci. Technol., 2 (2): 1 – 12.
- Havling, L.J, Beaton, D.J., Tisdale, L.S and Nelson, L.W (2006). Soil fertility and fertilizers: An introduction to nutrition management. Prentice – Hall of Indian, Private Limited. 7<sup>th</sup> edition. Pp 99 – 102.
- Horsfall, M. Jr and Spiff, I.A (2005). Equilibrium sorption study of Al<sup>3+</sup>, CO<sup>2+</sup> and Ag<sup>+</sup> in aqueous solution by Fluted Pumpkin (*Telfaria occidentalis* Hook F) Waste Biomass. Acta Chim Slov.
- Ikediobi, C.O., Onyia, G.O.C and Eluwah, C.E (1980). A rapid and inexpensive non-enzymatic assay for total cyanide in cassava (*Manihot esculenta crantz*) and cassava product. Agric. Biol. Chem., 44: 2803 – 2808.
- Jones, R.J and Ford, C.W (1972). Some factors affecting the oxalate content of the tropical glass (*Setaria sphacelata*). Aust. J. Exp. Agric., 12 (57): 400 – 406.
- Juo, A.S.R (1979). Selected methods of soils and plants analysis: Farming systems program – Manual series No. 1. Ibadan, IITA.
- Kansal, B.D, Bajaj, K.L and Kaaur, G (2005). Effect of different levels of Nitrogen and farmyard manure on yield and quality of spinach (*Spinace oleracea*). Plant J Plant Foods for Human Nutr., 31 (2); 163 – 170.
- Kriedenman, P.E (1964). Cyanide formation in sorghum alum in relation to nitrogen and phosphorus nutrition. Aust. J. Exp. Agric., 4 (12): 15 – 16.
- Lisiewska, Z., Kmiecik, W and Gebczynski, P (2006). Effect on mineral content of different methods of preparing frozen root vegetables. Food Sci. Technol. Int., 12 (6): 497 – 503.

- Mozafar, A (2005). Decreasing of NO<sub>3</sub> and increasing the Vitamin C contents spinach by a Nitrogen deprivation method. *J. Plant Foods for Human Nutr.*, 49 (2): 155 – 162.
- Muramoto, J (1999). Comparison of nitrate content in leafy vegetables from organic and conventional farms in California. Center for Agroecology and Sustainable Food System. University of California, Santa Cruz., 3; 40.
- Musa, A., Ezenwa, M.I.S., Oladiran, J.A., Akanya, H.O and Ogbadoyi, E.O (2010) Effect of soil nitrogen levels on some micronutrients, antinutrients and toxic substances in *Corchorus olitorius* grown in Minna, Nigeria. *Afr. J. Agric. Res.*, 5 (22): 3075 – 3081.
- Musa, A., Ogbadoyi, E.O., Oladiran, J.A., Ezenwa, M.I.S and Akanya, H.O (2011) Effect of fruiting on micronutrients, antinutrients and toxic substances in *Telfairia occidentalis* grown in Minna, Niger State, Nigeria. *Afr. J. environmental Sci. and Technol.*, 5(9): 710 – 716.
- Nkang, A., Omokaro, D., Egbe, A and Amanke, G (2003). Variations in fatty acid proportions during desiccation of *Telfairia Occidentalis* seeds harvested at physiological and agronomic maturity. *Afr. J. Biotechnol.*, 2 (2): 33 – 39.
- Ogbadoyi, E.O., Musa, A., Oladiran, J.A., Ezenwa, M.I.S and Akanya, H.O (2011). Antinutrient and micronutrients contents in leaves of *Telfairia occidentalis* Hook. f.: The effect of cooking and sun drying. *South Asian J. Exp. Biol.*, 1 (2): 61 – 68.
- Ojeniyi, S.O and Adeniyi, N.O (1999). Effect of poultry manure and NPK on soil fertility, nutrients and yield of maize at Akure South West Nigeria. Soil Science Society of Nigeria: Proceedings of the 25<sup>th</sup> Annual Conference. Pp 185 – 191.
- Ojiaka, O.A and Igwe, CU (2008). The nutritive, anti-nutritive and hepatotoxic properties of *Trichosanthes anguina* (Snake tomato) Fruits from Nigeria. *Pak. J. Nutri.*, 7 (1): 85 – 89.
- Oke, L.O (1966). Composition of some Nigeria leafy vegetables. *J. Am. Dietetic Assoc.*, 53: 130 – 132.
- Okoli, B.E and Nyanayo, B. L (1988). Polynology of *Telfairia L* (Cucurbitaceae). *Folia Geobotanica phytotaxonomica*. 23: 281 – 286.
- Oladele, S.B., Ayo, J.A and Aduadi, A.O (1997). The emergence of nitrate and nitrite poisoning in humans and domestic animals. *West Afr. J. Pharmacol. Drug Res.*, 13 (1 and 2): 50 – 58.
- Osunde, A.O and Alkassoum, A (1998). Growth, natural nodulation, and nutrient yield of selected multipurpose tree species in a Nigerian moist savana soil. *Agro. Afr. Numero Special*. 1: 361 – 373.
- Peter, K.B and Birger, L. M (2002). Dhurrin synthesis in sorghum is related at the transcriptional level and induced by nitrogen fertilization in order plants. *Plants physiol.*, 129: 1222 – 1231.
- Richard, D.W (1991). Cooperative Extension Service: Cooperative Extension work acts may 8 and June 30, 1914, as amended, Kansas State University, County Extension Councils. Extension Districts and U.S. Department of Agriculture Cooperating.
- Rickman, J.C., Bruhn, M.C and Barret, D.M (2007). Nutritional comparison of fresh, frozen and canned fruits and vegetables ii. Vitamin A and carotenoids, Vitamin E, minerals and fiber. *J. Sci. Food Agric.*, 88; 1185 – 1196.
- Rolinda, L.T and Ma, T.P.L (2008). Cyanide content of cassava cultivars at different fertility levels and stages of maturity. [Http://region10.dost.gov.ph/index.php?option=com-content&task=view&id=65&Itemid=77](http://region10.dost.gov.ph/index.php?option=com-content&task=view&id=65&Itemid=77).
- Safaa, A.M and Abd El Fattah, M.S (2007). Effect of nitrogen forms on nitrate contents and mineral composition of lettuce plants in sandy and calcareous soils. *J. Appl. Sci. Res.*, 3 (11): 1630 – 1636.
- Schippers, R.R (2000). African indigenous vegetables: An overview of the cultivated species. University Greenwich, England.
- Singh, P.P (2005). Influence of light intensity, fertilizers and Salinity on oxalate and mineral concentration of two vegetables (*Chenopodium Album L.* and *Chemopodium amaranthicolor L.*) *J. Plant Foods Nutr.* Springer Nether lands. 24 (2): 115 – 125.
- Sjoberg, A.M.K and Alanko, T. A (1994). Spectrophotometric determination of nitrate in baby food: Collaborative study. *J AOAC Int.*, 77 (2): 425–430.

- Sydenham, D (1985). Success in vegetable production. 1<sup>st</sup> edition. Macmillan London. Pp 23.
- Taiz, L., Zeiger, E (2002). Plant physiology. Sinauer Associates, Inc. Sunderland. Massachusetts. 3<sup>rd</sup> edition. Pp 370 – 372
- Tarfa, B.D., Uyovbisere, E.O., Chude, V.O., Raji, B.A and Yaro, D.T (2001). Effect of the complementary use of foliage of *Azadirachta indica*, *Parkia biglobosa* and NPK on yield and nutrients uptake of maize in a savana soil. Nig. J. Res., 2: 43 – 50.
- Virginia, W (2001). Nutritional quality of organic versus conventional fruits, vegetables and grains. J Vegetables Complement Med., 7 (2): 161 – 173.
- Waclaw, M and Stefan, S (2004). Effect of culinary processes on the content of nitrates and nitrites in potatoes. Pak. J. Nutr., 3 (6): 357 – 361.
- Yang, Y.J (1992). Effect of storage treatment on NO<sub>3</sub> and NO<sub>2</sub> contents in vegetables. J. Kor. Soc. Hortic. Sci., 33: 125 – 130.