



Received: 17th April-2013

Revised: 25th April -2013

Accepted: 26th April -2013

Research article

THE APPLICATION OF PRESERVATIVES ON QUALITY OF GLADIOLUS (GLADIOLUS GRANDIFLORUS) CUT SPIKE.

Amit K Awasthi^{a1}, P.K.Karhana^b and K.K.Pandey^c

^aSRF (Horticulture), VCSG College of Horticulture, Uttarakhand University of Horticulture and Forestry, Bharsar, Pauri-Garhwal, (Uttarakhand), INDIA.

^bDepartment of Horticulture RB (PG) College, Agra (UP) INDIA.

^cDepartment of Agriculture Statistics, VCSG college of Horticulture, Uttarakhand University of Horticulture and Forestry, Bharsar, Pauri-Garhwal, (Uttarakhand), INDIA

*Corresponding Author e-mail:awasthia83@gmail.com

ABSTRACT: In the present study attempts has been made to effect of two preservatives along with their chemical combination on gladiolus for western Uttar Pradesh. The lab of department of horticulture in RB (PG) College, Agra (UP) has been used for the experimentation in 2011. The CRD with four treatments and three replications has been used under the study. The preservatives has been taken AgNO₃ (25ppm), sucrose 2% AgNO₃ (25 ppm) + sucrose 2% and distil water (control) denoted as T1, T2, T3 and T4 respectively, for flower treatment. The result indicates that T1 taken minimum day for basal floret open followed by T3. Floret open (%) has been found maximum in T1 followed by T3. The T1 indicates as best preservatives in diameter of florets and fresh weight of flower. Our conclusion has been drawn that the T1 found as a best preservatives followed by T3 for the western U.P. during the season.

Key Words: AgNO₃, Gladiolus, Preservatives and Sucrose, cut spike.

INTRODUCTION:

In India, most of the hybrids were also developed by the Raja Bhadri, Governor of Himachal Pradesh from 1956-63 by collecting open pollinated seeds from different varieties recommended for that state. Similarly in Kartain (Kullu Valley), the work on assessment was also taken up from 1962. The cultivars were also collected from India and abroad at IARI, New Delhi. These were assessed and grown under plain conditions. Later the work was also started at IIHR at Hassaraghatta Bangalore. Similarly planned breeding was started in 1970 on wards, New Delhi, and later on IIHR, Bangalore, National Botanical Research Institute (N.B.R.I.) Lucknow (U.P.), Horticulture experiment and Training centre, Chaubattia, Ranikhet (Uttarakhand) etc. In northern plains of U.P. its commercial cultivation is gaining popularity due to prevailing congenial climatic conditions for planting from August to December and spikes are available from 1st week of November to end of April. In India most of the cultivars of the gladiolus were introduced from abroad except a few like Ratna's Butterfly, which were developed naturally by a nursery in Kalimpong. The use of floral preservatives is the most economical practical methods for extending post harvest life of gladiolus cut flower. The vase life of cut flower is influenced by constant water supply, checking of microbial growth prevention of ethylene formation and energy source. Several types of floral preservatives in form of germicides, ethylene antagonistic and source of energy (Sucrose) are in use to preserve the flower quality and extending post harvest longevity of cut flowers [8]. A lot of works have been done for the study of Gladiolus with the different chemicals in different part of India, but no work has been done in this direction for the Western Uttar Pradesh for Gladiolus. In the present study, an attempt has been made for the chemical effect as a growth regulator on Gladiolus cut spike for Agra district.

MATERIALS AND METHODS

The present study has been carried out in department of horticulture laboratory of the R.B.(PG) College, Agra (U.P.) during the February 2011. The experimental flowers were held in the laboratory at about $22 \pm 2^\circ\text{C}$ ambient room temperature and $80 \pm 5\%$ percent relative humidity (RH). Under the study 3 replication for each 4 treatments like T1, T2, T3 and T4 has been taken as AgNO_3 (25ppm), Sucrose (2%), AgNO_3 (25ppm) + Sucrose(2%) and Distilled water (control) respectively, under layout of CRD. Total 12 treatments and 2 flower scales per treatment has been taken for the study. The cut spikes have been harvested in the morning between 8 to 9 am. 5 gram of HgCl_2 was dissolved in 500 ml distilled water for disinfectant. Thereafter, in laboratory the spikes have cut to a constant length of 45 cm each and the fresh weight of cut spike for each treatment and replication has been recorded.

Preparation of Floral Preservative Solution

Two chemicals viz. AgNO_3 and Sucrose have used alone and combination of sucrose and AgNO_3 for quality parameter (preservatives) of cut gladiolus flower. The stock solution of silver nitrate (AgNO_3) and sucrose made on 1000 ppm prepared by dissolving proportionate weight in gram of individual chemical in distilled water and then final volume make up to 500 ml.

All the holding solution has been store in 500 ml flask. The cotton plug has putted in the 500 ml flask to avoid contamination, evaporation for further use formula used $01 \text{ f} = 1000 \text{ ppm} = 1000 \text{ mg/l}$.

Days to Basal Florets Open

Days to basal florets open has been recorded from date of placing to spike in holding solution to complete opening of basal florets.

Percentage of Florets Opened

The total numbers of florets on each spike have counted on the day of harvest and the total numbers of fully opened florets on each spike have counted florets.

$$\text{Percent of opened florets/spike} = \frac{\text{No. of Opened florets / spike}}{\text{Total number of florets / spike}}$$

Flower Diameter

Diameter of the floret has been measured on two perpendicular axis of the first, third and last fully opened pair of florets.

Fresh Weight change

Weight of spike has been measured on the day of experimental setup and at senescence. The difference between the weight of the container + solution + flower and the weight of container + solution recorded at every alternate day to measure the fresh weight change of flower during that particular period duration of time. The weight of the flower stems on the first day of each experiment has assumed to be 100 per cent. Subsequent weights have referred to as percentage of the initial value.

Statistical Analysis

The data were statistically analyzed with the help of computer using 'completely randomized design'. The treatments were compared with the help of critical difference suggested by [3] at 5% level significantly and the results thus obtained are presented with the help to tables.

$$\text{C.D.} = \sqrt{\frac{2\text{SE}}{r}} \times t_{\text{at } 5\%}$$

Where:

C.D. = Critical Difference

SE = Standard Error.

r = number of total treatments.

t at 5% = t value of error degree of freedom at 5% level of significant.

RESULTS

The result pertaining to the number of days required for first flower opening under the influence of various treatments has been presented in Table-1. The results shows that number of days required for first flower opening has been lowest in T1 (AgNO₃ 25 ppm) being 2.35 days followed by T3 (AgNO₃25ppm+ sucrose 2%)3.25 days. The result of present study presented has show maximum fully opened florets per spike were recorded in Sucrose 2% (92.10) and followed by AgNO₃ (25 ppm) (91.15). However, the minimum percentage of fully opened floret have recorded (80.56) in control (Distilled water), (Table-2). All the records three cases first fully opened, third fully opened & last fully opened. The AgNO₃ (25ppm) found best as 9.20 cm, 9.20 cm, 9.25 cm respectively. Followed by T2 (sucrose 2%) as 8.4cm, 8.12cm and 7.8cm respectively, (Table-3). The data related to fresh weight of spike has shown the weight has taken at experiment set up. The observation shows that which are deep in silver nitrate (AgNO₃) and sucrose. Solutions have gain weight the holing solution containing the AgNO₃ (25ppm) was being (36.17) and followed by control and rest have low performance of flower after harvest, (Table-4).

Table-1: Effect of various treatments on the days of basal floret open

Treatment	Days taken to basal floret open
T1	2.35
T2	3.82
T3	3.25
T4	2.74
SEm±	0.349
CD at 5%	1.101

Table-2: Effect of different treatment on open Florets in percentage

Treatments		Opened florets (%)
AgNO ₃ 25 PPM	T1	91.15 ± 4.78
Sucrose 2%	T3	92.10 ± 1.37
AgNO ₃ 25 PPM + Sucrose 2%	T3	83.51 ± 2.72
Distilled Water	T4	80.56 ± 3.44
SEm±		0.182
CD at 5%		0.622

Table-3: Effect of various treatments on diameter of different fully opened Pair.

Treatments	First fully opened florets Diameter (cm)	Third fully opened florets Diameter (cm)	Last fully opened florets Diameter (cm)
T1	9.20	9.12	9.25
T2	8.40	8.12	7.83
T3	7.80	9.74	9.23
T4	7.20	7.01	6.23
SEm±		0.121	
CD at 5%		0.442	

Table-4: Effect of various treatments on fresh weight of spike

Treatment	Spike weight at harvest	Flower weight after vase life	Loss weight of spike
T1	74.33	38.16	36.17
T2	69.50	56.66	12.84
T3	76.00	60.83	15.17
T4	73.33	49.50	23.83
SEm±	1.419	1.159	4.818
CD at 5%	2.89	3.42	14.40

DISCUSSION

Almost similar result has been found by [6] reported that fresh mass and volume of water uptake in spike were improved with sucrose application to vase solution. However, a significant increase in fresh weight was observed due to holding solution containing sucrose 5%+*HQC 200ppm in Cv. 'Hunting Song', 'Song', and "Spike span"[9]. [5] reported the maximum increase in fresh weight was also reported by Al₂(SO₄)₃ [4] and cobalt [7]; [12]. The basic requirement to maintain growth area preservations fulfills by these two biocides, firstly by supplying the tissue with carbohydrate and secondly by decreasing the water uptake of spike. Which accelerate floret opening. [11] reported that gladiolus spike held in vase solution containing 5% sucrose + AgNO₃ (50 ppm) + 8-HQC (300 ppm) + acidifier had highest fully opened florets as compared to other treatments and control. According to [1], GA-3150 ppm has found a good growth regulator for Akola in Maharastra during ravi season. [2], reported that different solution acid and Benzyladenin was a good growth regulator for flowering in lily for Rasht. [10] reported that the effect of different growth regulator in different solution for Tamilnadu, India.

CONCLUSION

The maximum day taken for opening of Basal floret has been recorded in AgNO₃ 25 ppm followed by sucrose 2%. The maximum floret opening has recorded under the treatment sucrose (2%) and followed by AgNO₃ (25ppm) and minimum under control (Distilled water). However, the maximum floret size of first third and last fully opened pair has been recorded under the treatment T1 (AgNO₃ 25 ppm) and followed by control (distilled water). All floral preservatives significantly affected the fresh weight. Among all the floral preservatives tested maximum and minimum loss in fresh weight has noted with treatments T1 and T3, respectively. The cut gladiolus treated with AgNO₃ and sucrose. However, the effect of preservatives like AgNO₃ is best for the future case study on gladiolus, in Western U.P.

REFERENCES

- [1] Chopde , N., Gonge, V.S., Nare, P.K. 2011. Effect of growth regulators on growth and flowering of Gladiolus, Asian Journal of Horticulture, 6(2), pp. 398-401.
- [2] Emami, H., Saeidina, M., Hatamzadeh, Bakhshi, D., Ghorbani, E. 2011. The Effect of Gibberellic acid and benzyladenine in growth and flowering of Lily (*lilium longiflorum*), Advances in Environmental Biology, 5(7) pp. 1606-1611.
- [3] Fisher R.A. 1924. The Influence of Rainfall on the Yield of London Wheat at Rothamsted, Phil. Trans. Roy. Soc., 213, pp. 89-142.
- [4] Gowda, J.V.N and Murthy, G.M.A., 1993. Effect of aluminum, calcium and sucrose on post harvest life of Gladiolus. In J. Prakash (ed.) Floriculture: Technology. Tracts and Trends, Oxford and IBH Pub Co Pvt. Ltd., New Delhi, pp. 492-495.
- [5] Marousky, F.J., 1969. Conditioning gladiolus spikes to maintenance of fresh weight with pretreatments of 8-HQC plus sucrose. Proc. Fla. State Hort. Soc. 82, pp. 411-414.
- [6] Mishra, R. L. 1996. Commercial production of Gladiolus. Floriculture Today, 1(2), pp.15-17.
- [7] Murali, T.P., Reddy, T.V., 1993b. Post harvest life gladiolus as influenced by sucrose and metal salts. Acta Hort. 343, pp. 313-320.
- [8] Shukla, R., Kher, M.A., 1979. A note on prolonging vase life of bougainvillea. Indian J. Hort., 36, pp. 93-95.
- [9] Song, C.U., Shin, D.G., Noor, I.S., Lee, J.S., 1992b. Studies on the vase life extension of cut gladiolus. J. Korean. Soc. Hort. Sci. 33(1), pp. 95-101.
- [10] Sudhakar, M., Kumar, S.R., 2012. Effect of growth regulators on growth, flowering and corm production of gladiolus (*gladiolus grandiflorus* L.), Ind. Jr. plant. Sci., 1(2-3), pp. 133-136.
- [11] Wang, X.D., Gu, Y.M., 1985. Effect of different solution of water equilibrium and maintenance of freshness of gladiolus hybridus. Hort Sangahi Agric Sci. and Technology. 6, pp. 16-17.
- [12] Zhou, W.Z., Wang, Y., Zhang, C.L., 1994. Experiment on a preservative for flower of vase gladiolus (*Gladiolus Hybridus*) Hort. J. Jiangru Agriculture College, 16(4) 27-30.